

Equine GM-CSF Do-It-Yourself ELISpot



Catalog Number: SPOT2424E-004
Storage: 2-8°C
Stability: 12 months (from date of receipt)
Country of Origin: USA

Description: Contains Capture Antibody, Standard Control, and Detection Antibody for development of up to two (2) ELISpot assays. Other required materials and reagents are not supplied. The antibodies have been determined to function in an ELISpot with the Standard Control provided. Components may also be purchased separately.

Included Components:	Description	Usage	Quantity	Catalog Number
	Anti-Equine GM-CSF Polyclonal Antibody	Capture Antibody	100 µg	KP1926E-100
	Recombinant Equine GM-CSF	Standard Control	5 µg	RP0022E-005
	Biotinylated Anti-Equine GM-CSF Polyclonal Antibody	Detection Antibody	50 µg x 2	KPB1927E-050

Additional Materials and Reagents Required:	Material/Reagent	Suggested Formulation
	96-well ELISpot Plate	96-well, sterile, filter membrane ELISpot plate <i>ELISpot Plate: Catalog # AR2307-001</i>
	Sterile DPBS	0.008M sodium phosphate, 0.002M potassium phosphate, 0.14M sodium chloride, 0.01M potassium chloride, pH 7.4, sterile filtered
	Complete Medium/ Blocking Buffer	Sterile medium containing 10% heat-inactivated Fetal Bovine Serum (FBS) or Equine Serum (ES) in desired cell culture medium, such as RPMI. If required, serum-free medium for cell culture can also be used.
	Reagent Diluent	4% BSA in DPBS, 0.2 µm filtered
	Wash Buffer	0.05% Tween®-20 in DPBS
	Streptavidin-AP	Enzymatic reagent to react with biotinylated detection antibody <i>Streptavidin-AP: Catalog # AR2308-001</i>
	BCIP/NBT Substrate	5-bromo-4-chloro-3-indolylphosphate and nitroblue tetrazolium substrate to react with AP. <i>BCIP/NBT Substrate: Catalog # AR2309-025</i>

Time Requirements:		
	1 overnight incubation	Coated plate preparation
	1-2 overnight incubation	Cell stimulation as desired
	2.25+ hours	Assay development



Reagents for Animal Model and Animal Health Research

1000 Westgate Drive, St. Paul, MN 55114
 Customer.Service@KingfisherBiotech.com
 KingfisherBiotech.com
 651.646.0089

Do-It-Yourself ELISpot

Suggested ELISpot Controls:

Control	Suggested Formulation
Standard Control	Recombinant Protein at 50 ng/mL in Complete Medium.
Blank Control	Complete Medium only.
Positive Control	Cells incubated with a polyclonal activator such as PHA, PMA/Ionomycin, ConA, or other, as desired.
Negative Control	Cells in Complete Medium only.

ELISpot Protocol: Sterile Conditions Suggested:

1. Prepare Capture Antibody Working Solution in DPBS at 4 µg/mL, or as desired.
2. Add 100 µL of Capture Antibody Working Solution to each well.
3. Cover plate and incubate at room temperature (20-25°C) for 12-24 hours.
4. Empty Capture Antibody Working Solution from each well.
5. Add 250 µL of Complete Medium/Blocking Buffer to each well.
6. Cover plate and incubate at room temperature for 1-3 hours.
Tip: Prepare controls and samples for ELISpot during Blocking Buffer incubation.
7. Empty Blocking Buffer from each well.
8. Add 100 µL of controls or samples to appropriate wells.
Note: Run each control or sample in triplicate.
9. Cover plate and incubate at 37°C, 5% CO₂ for 12-48 hours, as desired.
DO NOT DISTURB PLATE DURING INCUBATION!

No Sterile Conditions Required:

10. Wash plate SIX times with Wash Buffer.
Note: Vigorous plate washing is essential; however, care should be taken to avoid puncturing the filter membrane ELISpot plate during wash steps. If using an automatic plate washer, confirm and adjust the height of the manifold dispenser prior to use. Take care to avoid microbial contamination of equipment. Automated plate washers can easily become contaminated thereby causing assay variability.
11. Prepare Detection Antibody Working Solution in Reagent Diluent at 4 µg/mL, or as desired.
12. Add 100 µL of Detection Antibody Working Solution to each well.
13. Cover plate and incubate at room temperature for 1 hour.
14. Wash plate SIX times with Wash Buffer as described in step 10.
15. Prepare Streptavidin-AP in Reagent Diluent at desired working concentration.
16. Add 100 µL of Streptavidin-AP Working Solution to each well.
17. Cover plate and incubate at room temperature for 1 hour.
18. Wash plate SIX times with Wash Buffer as described in step 10.
19. Add 100 µL of BCIP/NBT Substrate Solution to each well.
20. Develop the plate at room temperature for up to 15 minutes.
21. Stop reaction by flooding the plate with ultrapure water. Remove under tray and flood bottom of tray with ultrapure water.
22. Dry plate and evaluate results.

Warranty:

Products are warranted by Kingfisher Biotech, Inc. to meet stated product specifications and to conform to label descriptions when used, handled and stored according to instructions. Unless otherwise stated, this warranty is limited to one year from date of sale or expiry date, whichever comes first. Kingfisher Biotech's sole liability for the product is limited to replacement of the product or refund of the purchase price. Kingfisher Biotech products are supplied for research applications. They are not intended for medicinal, diagnostic or therapeutic use. The products may not be resold, modified for resale or used to manufacture commercial products without prior written approval from Kingfisher Biotech, Inc.

Tween is a registered trademark of ICI Americas.
© 08/2024



Reagents for Animal Model and Animal Health Research

1000 Westgate Drive, St. Paul, MN 55114
Customer.Service@KingfisherBiotech.com
KingfisherBiotech.com
651.646.0089